




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 9/00, 31/66, 31/70 A61K 35/78, C07D 207/00 C07D 327/00, 333/02, 335/00 C07F 9/02, C07H 15/00, 17/00		A1	(11) International Publication Number: WO 94/08563 (43) International Publication Date: 28 April 1994 (28.04.94)
(21) International Application Number: PCT/US93/09623 (22) International Filing Date: 8 October 1993 (08.10.93) (30) Priority data: 958,416 8 October 1992 (08.10.92) US (60) Parent Application or Grant (63) Related by Continuation US 958,416 (CIP) Filed on 8 October 1992 (08.10.92) (71) Applicant (for all designated States except US): SHAMAN PHARMACEUTICAL, INC. [US/US]; 887 Industrial Road, San Carlos, CA 94070 (US). (72) Inventors; and (75) Inventors/Applicants (for US only) : TEMPESTA, Michael [US/US]; 854 Tierra Alta, Moss Beach, CA 94038 (US). JOLAD, Shivanand, D. [IN/US]; 1710 San Carlos Ave- nue, #1, San Carlos, CA 94070 (US). KING, Steven [US/US]; 201 Wienke Way, Moss Beach, CA 94038 (US). MAO, Guohua [CN/US]; 137 Elm Street, San Car- los, CA 94070 (US). BRUENING, Reimar, C. [DE/US]; 3328 Brittan Avenue, #12, San Carlos, CA 94070 (US). KUO, John, E. [US/US]; 937 Jackson Street, Mountain View, CA 94043 (US). TRUONG, Thien, V. [US/US]; 6400 Christie Avenue, #3103, Emeryville, CA 94608 (US). BIERER, Donald, J. [US/US]; 880 Campus Drive, #122, Daly City, CA 94015 (US). DENER, Jeffrey, M. [US/US]; 850 Campus Drive, #106, Daly City, CA 94015 (US).		(74) Agents: BALDWIN, Geraldine, F. et al.; Pennie & Ed- monds, 1155 Avenue of the Americas, New York, NY 10036 (US). (81) Designated States: AU, BB, BG, BR, BY, CA, CZ, FI, HU, JP, KR, KZ, LK, LV, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.	

(54) Title: NOVEL CLASS OF PHOSPHOCHOLINE DERIVATIVES HAVING ANTIFUNGAL ACTIVITY**(57) Abstract**

Certain phosphocholine derivatives having substantial antifungal therapeutic activity are disclosed. The phosphocholine derivatives may be chemically synthesized, enzymatically prepared or extracted from the plant *Iribachia alata*. The phosphocholine derivatives are useful in treating fungal infections including those which are dermatophytic, systemic, ophthalmic and vaginal.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	CN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CF	Central African Republic	KR	Republic of Korea	SD	Sudan
CG	Congo	KZ	Kazakhstan	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovak Republic
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MG	Madagascar	UA	Ukraine
DE	Germany	ML	Mali	US	United States of America
DK	Denmark	MN	Mongolia	UZ	Uzbekistan
ES	Spain			VN	Viet Nam
FI	Finland				

- 1 -

NOVEL CLASS OF PHOSPHOCHOLINE DERIVATIVES
HAVING ANTIFUNGAL ACTIVITY

5 This is a continuation-in-part application of
U.S. patent application Serial No. 07/958,416, filed
October 8, 1993, the entire disclosure of which is
incorporated by reference.

10 1. Field of the Invention

 This invention relates to new classes of
phosphocholine derivatives as well as to various
methods for preparing these compounds -- including
synthetic, enzymatic and extractive using certain
15 plants. The phosphocholine derivatives of the
invention are non-toxic and exhibit substantial
antifungal activity in slowing fungal growth and in
killing fungi.

20 2. Background of the Invention

 The plant species *Irlbachia alata* has been used
as an anti-infective agent in the Peruvian Amazon
region. The leaves are squeezed and the liquid is
applied to infected skin sores. The same liquid from
25 the leaves is applied to skin problems and skin fungal
infections. It is utilized to treat vaginal yeast
infections.

Irlbachia alata is one species of 10-12 species
of the plant family Gentianaceae. These species occur
30 in tropical South America especially in the Amazon and
Negro River basins. The plants in the genus *Irlbachia*
are generally low herbs characteristically with 3-5
plinerved leaves. The most consistent diagnostic
feature for the genus is the pollen morphology.

35 A reference to *Irlbachia alata* and related
species was made in 1775 by the French scientist Fusee

Aublet (Aublet, F. 1775, Histoire des Plantes de la
Guiane Francoise, Didot, Paris). The ethnobotanical
notes from this reference were subsequently compiled
5 and republished in English. Aublet noted the
following about two species in the genus *Irlbachia*:

Irlbachia alata The entire plant is bitter.
It is used to clear obstructions; I (Aublet) have used
it with good results. The species is called "Bois
10 creux" (Hollow wood) by the Creoles.

Irlbachia pururascens All parts of this plant
are bitter. It is used as an apertif and to reduce
fever.

15 3. Summary of the Invention


We have discovered a class of phosphocholine
derivatives (Class I) having extraordinary antifungal
activity.

Structurally, these compounds are phosphocholine
20 derivatives (1 or 2-deacyl-phosphatidyl cholines) in
which the 1 or 2-OH-group of the glycerol moiety has
been glycosylated with glucose, galactose, arabinose,
mannose, rhamnose or another sugar. The basic
chemical structure may be drawn as follows:

25

30

35




wherein one of R or R' is a sugar moiety and the other is an acyl or sugar moiety.

15 The molecular backbone common to all members of this class of compounds is drawn above. The acyl-group can be any long-chain fatty acid, while the sugar unit can be any of the sugars commonly found in plants, including but not limited to glucose,
20 galactose, arabinose, mannose, rhamnose, or another naturally occurring sugar.

We have additionally found a structurally related class of phosphocholine derivatives of similar or greater antifungal activity than the above-discussed class of phosphocholine derivatives (i.e., Class I).

25 One novel class of phosphocholine derivatives (Class II) having antifungal activity has the basic structure shown below:



where Q is C2 to C30 alkyl, alkenyl, alkynyl, branched alkyl, branched alkenyl, or branched alkynyl;

Z is oxygen or sulfur; X and Y are independent oxygen, sulfur, CH_2 , CF_2 , or N-R_1 ;


A, B, and T are independently alkyl, alkenyl, alkynyl, branched alkyl, branched alkenyl, or branched alkynyl radicals of C1 to C20 chain lengths; are independently or together cycloalkyl or bridged cycloalkyl radicals of ring size C3 to C20, or cycloalkenyl, bridged cycloalkenyl or cyclo(polyene)radicals of ring size C4 to C20, cycloalkynyl, bridged cycloalkynyl or cyclo(polyalkynyl)radicals of ring size C8 to C20;

D is oxygen, sulfur, CH_2 , CF_2 , or N-R_2 ;

F is alkyl, alkenyl, alkynyl, branched alkyl, branched alkenyl, branched alkynyl, cycloalkyl, bridged cycloalkyl, cycloalkenyl or cycloalkynyl radicals containing C1 to C20 carbon atoms;

R_1 and R_2 are independently hydrogen, alkyl, alkenyl, alkynyl, branched alkyl, branched alkenyl, branched alkynyl, cycloalkyl, bridged cycloalkyl, cycloalkenyl, bridged cycloalkenyl or cycloalkynyl radicals containing C1 to C20 carbon atoms, or any protecting group described in the book "Protecting Groups in Organic Synthesis" by Theodora Greene and Peter G.M. Wuts.

Another class of phosphocholine derivatives (Class III) having antifungal activity has the following structures:



35


where AA, BB, DD are independent of each other, equal to each other, or interchanged as shown above, the central carbon atom can be either the R and S
 5 optical stereoisomer or a mixture of R and S stereoisomers, and where AA, BB, and CC are defined as follows:

where AA, is A-J with A being attached to the carbon atom of the three carbon central unit and J is
 10 defined below;

BB is B-Y, with B being attached to the carbon atom of the three carbon central unit and Y is defined below:

DD is

15



20 where W, E, G and Q are defined below;

A is oxygen, sulfur, CH₂, CF₂ or N-R₁;


B is oxygen, sulfur, CH₂, CF₂ or N-R₂;

D is oxygen, sulfur, CH₂, CF₂ or N-R₃;

Y is alkyl, alkenyl, alkynyl, poly(alkenyl),
 25 poly(alkynyl), or poly(alkenoalkynyl) radicals
 comprised of C1 to C20 carbon atoms chain lengths, or
 alkanoyl, alkenoyl, alkynoyl, poly(alken)oyl,
 poly(alkyn)oyl or poly(alkenoalkyn)oyl radicals
 comprised of C2 to C20 chain lengths or alkyloxy,
 30 alkenyloxy, alkynyloxy, poly(alkenyl)oxy,
 poly(alkynyl)oxy, poly(alkenoalkynyl)oxy radicals
 comprised of C1 to C20 carbon atoms;

35

J is a furanose or pyranose radical of the type:



- 10 where X is oxygen, sulfur, CH₂, CF₂ or N-R₄;
 F, K, L and M are independently hydrogen,
 hydroxyl, protected hydroxyl (as described in the book
 "Protecting Groups in Organic Synthesis" by Theodora
 Greene and Peter G.M. Wuts), alkyloxy, thiol,
 15 alkylthio, arylthio, alkylsulfonyl, arylsulfonyl,
 amino, ammonium, alkylamino, alkylammonium,
 dialkylamino, dialkylammonium, trialkylamino,
 trialkylammonium where the alkyl chain on nitrogen is
 comprised of C1 to C20 carbon atoms; or alkyl,
 20 alkenyl, or alkynyl radicals comprised of C1 to C20
 carbon atoms.

Z is oxygen or sulfur

E is oxygen, sulfur, CH₂, CF₂ or N-R₅;

- G is alkyl, branched alkyl, cycloalkyl or bridged
 25 cycloalkyl radicals of C1 to C20 chain lengths;


- Q is halogen, hydroxyl, protected hydroxyl
 utilizing any protecting groups described in the book
 "Protecting Groups in Organic Synthesis" by Theodora
 Greene and Peter G.M. Wuts, O-arylsulfonyl-, O-
 30 alkylsulfonyl- or O-(perfluoroalkyl)sulfonyloxy,
 amino, ammonium, alkylamino, alkylammonium,
 dialkylamino, dialkylammonium, trialkylamino,
 trialkylammonium where the alkyl chains on nitrogen
 are C1 to C20, or Q=NR₁R₂R₃, where R₁, R₂, or R₃ can
 35 independently or together be a mixture of alkyl groups
 of C1 to C20 in chain length and a protecting group

described in the book "Protecting Groups in Organic
Synthesis" by Theodora Green and Peter G.M. Wuts, and
R₁ can equal R₂, R₂ can equal R₃, or R₁ can equal R₃,
5 which can equal R₃;


R₁, R₂, R₃, R₄ and R₅ are independently alkyl,
alkenyl, alkynyl, branched alkyl, branched alkenyl,
branched alkynyl, cycloalkyl, bridged cycloalkyl,
cycloalkenyl or cycloalkynyl radicals of C1 to C20
10 chain lengths, or any protecting group described in
the book "Protecting Groups in Organic Synthesis" by
Theodora Greene and Peter G.M. Wuts;

where W₁ and W₂ are P(-OR) (with R being phenyl,
phenylmethyl, or negatively-charged oxygen), S=O,
15 carbon, or sulfur, provided that if W₁ is not P(-OR) W₂
is P(-OR) and provided that if J is a furanose or
pyranose radical then W₁ is P(-OR).

A preferred subgroup of the above-described
Class III of phosphocholine derivatives have the
20 following structures:






25



30

35



25 where R_1 is phenyl or phenylmethyl, hydrogen, or nil;

R_2 is hydrogen, phenylmethyl, or any protecting group described in the book "Protecting Group in Organic Synthesis" by Theodora Green and Peter G.M.

30 Wuts which can be cleaved by hydrogenolysis;

AA, BB, and Q are as defined above

where the central carbon atom of the three carbon unit is either the R optical isomer, the S optical isomer, or any mixture of the two optical isomers

35 thereof;

Another preferred subgroup of the above-described Class III of phosphocholine derivatives have the following structures:

5

10




15

20


25

30


35



5



10



15

where R_1 is phenyl or phenylmethyl, hydrogen, or nil; R_2 is hydrogen, phenyl methyl or any protecting group described in the book "Protecting Groups in Organic Synthesis" by Theodora Greene and Peter G.M.


20 Wuts which can be cleaved by hydrogenolysis;

R_3 is hydrogen or a protecting group as described in the book "Protecting Groups in Organic Synthesis" by Theodora Greene and Peter G.M. Wuts.;

25 where the central carbon atom of the three carbon unit is either the R optical isomer, the S optical isomer, or any mixture of the two optical isomers thereof; and Q is defined above.

Still another preferred subgroup of the above-described Class III of phosphocholine derivatives have
30 the following structures:

35



where R_1 is phenyl or phenylmethyl, hydrogen, or
 10 nil;

R_2 is a protecting group as described in the book
 "Protecting Groups in Organic Synthesis" by Theodora
 Greene and Peter G.M. Wuts, or hydrogen if R_1 is not
 hydrogen;

15 and Q is defined above.

We have further found a novel, generally
 applicable method for the synthesis of the above
 described broad classes of phosphocholine derivatives
 (Classes I, II and III).

20

4. Brief Description of the Drawings

Fig. 1 is the FTIR spectrum of the composition
 comprising a phosphocholine derivative obtained from
Irlbachia alata.

25 Fig. 2 is the proton NMR spectrum of the
 composition comprising a phosphocholine derivative
 obtained from *Irlbachia alata* in D_2O at 400 MHz.

Fig. 3 is the F $\bar{A}B$ /MB mass spectrum of the
 composition comprising a phosphocholine derivative
 30 obtained from *Irlbachia alata*.

5. Detailed Description of the Invention

The glycosylated lysolecithins of the invention
 can be prepared by synthetic methods or by enzymatic
 35 methods. The phosphocholine derivatives can be


prepared either by synthetic methods or by methods entailing extraction from plant materials.

5 5.1. Chemical Synthesis of phosphocholine derivatives

A wide variety of compounds having accessible alcoholic functionalities can be glycosylated following the classic Koenigs-Knorr methodology. Bochkov, A.F. and Zaikov. G.E., Chemistry of the O-Glycosidic Bond. Pergamon Press, 1979. As part of the synthetic route to phosphocholine derivatives with sugar, all but the anomeric hydroxyl group of the sugar to be introduced are protected either as esters or ethers, while the anomeric hydroxyl is being replaced by a halogen. The aglycon-sugar linkage is then formed via alcoholysis. Finally, the protective groups are selectively removed.

In the present invention, benzyl ethers or the benzilidene moiety are the preferred protecting group, since they can be selectively removed by catalytic hydrogenation, while leaving the sensitive acyl-glycerol linkage intact. The glycosidation requires silver, mercury (*Helferich* modification), or cadmium salts as catalytic halogen abstractor, in the presence of a dehydrating agent (Timell, T.E., *Can.J. Chem.* 1964, 42, 1456; Dejter-Juszynsky, M. and Flowers, H.M., *Carbohydr. Res.* 1973, 30, 287; Marousek, V., Lucas, T.J., Wheat, P.E., and Schuerch, C., *Carbohydr. Res.* 1978, 60, 85), and with or without auxiliaries such as crown-ethers. (Knöchel, A. Ger, R., and Thiem, J. *Tetrahedron Letters* 1974, 551) More recent methodology makes use of the halogen-abstracting power of non-nucleophilic bases such as diisopropylethylamine and/or of molecular sieves in an anhydrous media. (Garegg, P.J. and Norberg, T., *Carbohydr. Res.* 1976, 52, 235) The following

synthetic scheme is based on the latter reaction sequence:



The synthetic two-step scheme outlined above can be conducted with commercially available materials.

2,3,4,6-Tetrabenzyl- 2,3-dibenzyl-4,6-benzylidene-glucose can be converted into the 1-bromo- or 1-O-triflate compound by standard methodology. Leroux, J. and Perlin, A.S. *Carbohydr. Res.* 1976, 47, C8. The corresponding phosphocholine derivatives are available through AVANTI POLAR LIPIDS, Inc. All other reagents are available from ALDRICH. The methodology outlined above is also applicable to either 1-acyl or 2-acyl (1-acyl detailed above).

35




5.2. Enzymatic preparation of 1 or 2 glycosylated lysolecithins

As an alternative to the synthetic sequence outlined above, an *in vitro* enzymatic glycosidation simulating the biosynthetic process will produce the desired compounds in comparable yields. The natural glycosidation catalysts are glycosyltransferases. These enzymes operate with uridinediphospho-glycosides (UDP-sugars) as substrates and ATP as the energy source. While the enzymes have to be prepared from fresh plant material, UDP-sugars, ATP, as well as the respective phosphocholine derivatives are commercially available. This synthesis has the advantage of being essentially a one-step process with the high selectivity and yields expected from an enzymatic reaction. The following scheme describes the preparation of a glucoside. Other transferases, not specific to glucose, could be applied in the preparation of glycosylated lysolecithins with other sugars as well:

25

30

35



30

5.3. Total Synthesis of Phosphocholine Derivatives

A general synthetic method of synthesizing phosphocholine derivatives of the various structures described in section 3 is outlined as follows.

35 An alcohol is phosphorylated or glycosylated.
The product is subsequently deprotected. The

deprotected product is then alkylated or esterified to produce the phosphocholine derivatives. The general scheme for this outlined synthetic method is shown

5 below.

10

15


20

25

30

35

General Scheme for the Synthesis of Known and Novel Lysolecithins



Glycerol Derivative I can be either the R or S optical isomer, racemic, or a mixture of R and S isomers

⇒ Implies that a number of synthetic transformations are required

R₁ = Sugar, carbocyclic sugar, functionalized sugar derivative, etc.

R₂ = Phosphate or phosphate isostere moiety

R₃ = alkyl, alkanoyl, alkenyl, alkenoyl, etc.

X, Y, and Z can be C, O, N, S independently or equal to each other

5.4. Methods of Use

The phosphocholine derivative in Classes I, II and III are all useful in treating fungal infection by the administration to a warm-blooded animal of a therapeutically effective amount of a phosphocholine derivative. The pharmaceutical composition comprising the phosphocholine derivative used for such administration may also contain pharmaceutically acceptable excipients and carriers.

Phosphocholine derivatives in Classes I and II are believed to be novel compositions.

In order to treat a fungal infection, the antifungal agent of Classes I, II and III may be administered to a warm-blooded animal intravenously, intraperitoneally, subcutaneously, intramuscularly, orally, topically, by aerosol, or combinations thereof.

The antifungal agent of phosphocholine derivatives in Class II can be administered intravenously in a range of about 0.1 to about 10 mg/kg.

The fungal agent of Class II can be administered intraperitoneally in a range of about 0.1 to about 10 mg/kg.

The fungal agent of Class II can be administered subcutaneously in a range of about 1 to about 20.

The fungal agent of Class II can be administered intramuscularly in a range of about 1 to about 20.

The fungal agent of Class II can be administered orally in a range of about 5.0 to about 30 mg/kg.

The fungal agent of Class II can be administered topically in a range of about 5.0 to about 15% by weight.

The fungal agent of Class II can be administered by aerosol in a range of about 5.0 to about 30 mg/kg/day.

- 5 The above dosage ranges may need to be doubled for those phosphocholine derivatives in Class I and III with lower antifungal activity which are identical or similar to those in table 2 (see below).

10 6. Extraction of phosphocholine derivatives from plants

Plants are not known to contain phosphocholine derivatives.


- 15 The general manner of chemical extraction from the plants can be summarized as follows.

- 20 The plant source material, such as the whole plant, the roots, leaves, stem and/or latex of the plant, is extracted with water and/or a water miscible solvent. The preferred solvents are alcohol of 1-3 carbon atoms or acetone. The aqueous extract is extracted with butanol. The butanol-soluble fraction is subjected to gel filtration (e.g., over Sephadex), reversed-phase column chromatography (e.g., C-8), or gel-permeation chromatography (e.g., divinyl benzene cross-linked gels) such as PL-GEL or membranes (e.g., an Amicon membrane) using water or water and a water miscible solvent, with or without a buffer, as the mobile phase. The water miscible solvent is preferably a 1-3 carbon alcohol, acetone or
- 25 acetonitrile.
- 30

The useful phosphocholine derivatives containing compound is the fraction detected by NMR spectroscopy.

- 35 A specific member of the class of phosphocholine derivatives of the present invention is

2-palmitoyl-1-O-glycopyranosyllysolecithin shown
below:




2-Palmitoyl-1-O-glucopyranosyllysolecithin

15

We have found that 2-palmitoyl-1-O-
20 glucopyranosyllysolecithin is a relatively active
antifungal agent similar in activity to
L-a-Lysophosphatidyl inositol, discussed in Table 2
below.

We have found that one of the most active
25 antifungal compounds has the following structure.



1,22-docosan diol bisphosphocholine ester.

35

6.1. Extraction

We have isolated by chemical extraction 1,22-docosan diol biphosphocholine ester, the active
5 antifungal compound contained in the plant *Irlbachia alata*. The leaves of *Irlbachia alata* were milled and 200g of the milled leaves was extracted with 1L of dichloromethane/isopropanol (1:1 v/v) at room
10 temperature for 24 hours. The extracted material was separated from the marc (i.e., residual of the plant after solvent extraction) and discarded. The marc was then extracted with 1.5L of isopropanol/water (1:1 v/v) at room temperature for 24 hours. The marc was
15 separated from the extract and discarded. The isopropanol/water (1:1 v/v) soluble extract was partitioned between water and ethyl acetate. The ethyl acetate phase was separated and discarded. The water soluble phase, after extraction with n-butanol, was then discarded. The n-butanol phase was subjected
20 to filtration over two Sephadex LH-20 gel columns using 90% aqueous ethanol (for first filtration) and 20% aqueous acetone (for second filtration) as the mobile phases. 1,22-docosandiol biphosphocholine ester was collected from the early fractions of each
25 gel filtration.

We believe that several related genera are the same and/or closely related to the genus *Irlbachia*, and would have similar medicinal properties. One
30 species from a closely related genus, *Lisianthus nigrens* is used in Mexico. The leaves are applied as a poultice to treat fungal infections of the skin, feet, ankles and hands. A decoction of the root is also taken orally as a "bitter" and as a febrifuge. Another species *Lisianthus alatus* is considered to be
35 the same as *Irlbachia alata*. Another species and genus of interest is *Chelonanthus alatus*. There are

several uses described for *Chelonanthus alatus*, including oral decoctions to treat smallpox, fevers and for gastric disturbances.

5

6.2. Spectral Characteristics

The isolated phosphocholine derivative fraction containing 1,22-docosandiol bisphosphocholine ester has the characteristic IR, proton NMR and FAB⁺ mass spectra shown in Figs. 1, 2 and 3, respectively.

10

The IR spectrum has peaks at approximately 1060, 1220, 1475, 1600-1700, 2850, 2950 and 3400 cm⁻¹.

The ¹H NMR spectrum has major peaks at δ 1.2, 1.4, 1.7, 3.1, 3.5, 3.7 and 4.3.

15

The FAB⁺/MB mass spectrum has major peaks (>40%) at m/z 657, 612, 587, 586, 555, 493, 491, 475, 403, 277, 233, 201, 194, 179, 168, 165 and 163.

The high resolution mass spectrum (FAB⁺) has a molecular ion at 673.4669 amu.

20

6.3. Total Synthesis of 2-palmitoyl-1-O-glucopyranosyllysolecithin Experimental Section

General. Tetrahydrofuran (THF) was distilled from potassium/benzophenone; benzene, triethylamine, and methylene chloride, N-methylmorpholine, and benzyl alcohol were distilled from calcium hydride; 2-bromoethylphosphorodichloridate was prepared according to the procedure reported by Baumann et al Lipids, 17, 453 (1982) and was freshly distilled prior to use; trifluoromethanesulfonic anhydride was freshly distilled under inert atmosphere; O- α -D-(Glucopyranosyl)trichloroacetimidate was prepared by the method of Schmidt. (a) R. R. Schmidt, J. Michael, Angew. Chem. Int. Ed Engl. (1980), 19, 731; (b) R. R. Schmidt, J. Michael, Tetrahedron Lett. (1984), 25, 821. Anhydrous dimethylformamide (DMF) was obtained

35

from Aldrich. S-(+)-1,2-O-isopropylidene glycerol and R-(-)-1,2-O-isopropylidene glycerol were obtained from Lancaster. 2,3,4,6-Tetra-O-benzyl-D-glucopyranose was
5 obtained from Sigma. Preparative thin layer chromatography plates was performed on Whatman 2000 μ TLC silica gel plates. Flash column chromatography was performed on Whatman 230-400 mesh silica gel using nitrogen pressure. ^1H and ^{13}C NMR were provided by using
10 a Varian 400 MHz spectrometer with chloroform as an internal reference unless otherwise noted. NMR shifts were expressed in ppm downfield from internal tetramethylsilane. Carbon 13 multiplicities as determined by DEPT experiments are reported in
15 parentheses following the chemical shift value according to the following format: (0) for quaternary carbon, (1) for methine carbon, (2) for methylene carbon, and (3) for methyl carbons. NMR assignments were determined on the basis of COSY, HMQC, and HMBC
20 and DEPT experiments performed on selected intermediates. NMR coupling constants are reported in Hertz. Melting points were determined using a Buchi model 535 melting point apparatus and are uncorrected.


The synthetic routes for the total synthesis of
25 2-palmitoyl-1-O-glucopyranosyllysolecithin are outlined in the following diagrams and detailed in the subsequent discussion that refer to these diagrams.

30


35

- 24 -

Scheme 1. Synthesis of the (S) SP-19501: Preparation of the Re-isomeric Glycerol Alcohols




Scheme 2. Synthesis of the (R) SP-19501: Preparation of the Regioisomeric Glycerol Alcohols




- 26 -


Scheme 3. Synthesis of (S) SP-19501



- 27 -

Scheme 4. Synthesis of (R) SP-19501





(R) 2,3-O-Isopropylidene-1-O-trifluoromethylsulfonyl--
glycerol. A nitrogen-purged 250-mL three-necked
roundbottomed flask fitted with a thermometer,
5 stopper, and septum was charged with S-(+)-1,2-O-
isopropylidene glycerol (1.0 g, 7.6 mmol) dissolved in
benzene (75 ml). Triethylamine (1.25 mL, 9.0 mmol)
was injected into the solution, and the reaction
mixture was chilled until a cloudy solution appeared.
10 Trifluoromethanesulfonic anhydride (1.25 mL, 7.6 mmol)
was then added, and the reaction was stirred for 30
minutes with the temperature maintained at 5°C. The
solution was then filtered through a bed of silica.
The filtrate was concentrated under reduced pressure
15 at 30°C to give an orange/brown oil (1.84 g, 7.0 mmol)
in 92% yield which was used directly for the next
step.

(2R) [1-O-(2,3,4,,6-Tetra-O-benzyl- β -D-
20 glucopyranosyl)-2,3-O-isopropylidene] glycerol 1

2,3,4,6-Tetra-O-benzyl-D-glucopyranose (100 g,
0.182 mol) was dissolved in THF (1.4 L) and chilled to
-10°C in a nitrogen-purged 3-L three-necked morton
25 flask fitted with a thermometer, stopper, and
mechanical stirrer. Sodium hydride 60% in oil (16.1
g, -0.403 mol) was added in 4 increments over 10
minutes, and the solution was stirred for 30 minutes.

(R) 2,3-O-Isopropylidene-1-O-
30 trifluoromethylsulfonylglycerol (60.0 g, 0.227 mol)
dissolved in THF (500 mL) was then dropped via an
addition funnel into the reaction mixture over a 30
minute period. The solution was stirred at -10°C for
7 hours. Methanol (200 mL) was added dropwise to
35 quench excess sodium hydride, the resulting brown
solution was rotary evaporated under reduced pressure

and then the residue redissolved in chloroform (750 mL). The organic layer was washed with water (2 x 750 mL). The combined aqueous layers were washed with
5 chloroform (3 x 500 mL). Organic layers were pooled and rotary evaporated under reduced pressure to give a white solid which contained both α and β -epimers of the desired product. The solid was triturated with
10 diethyl ether to give a white solid of purely β -product and a mother liquor which contained α and β -epimers. The mother liquor was concentrated and flash chromatographed (silica gel, 20% ethyl acetate/hexane). Yield of the solid white β -epimer product (81 g, 0.123 mol) was 68%, mp 91-91.7°C (lit 83-84°C); $^1\text{H-NMR}$ (CDCl_3) δ 7.4-7.29 (m, 18H), 7.20 (m, 2H), 4.96 (d, 2H $J=10.8$), 4.84 (t, 2H, $J=10.8$), 4.75 (d, 1H, $J=10.8$), 4.65 (d, 1H, $J=12.4$), 4.6-4.54 (overlapping dd, 2H, $J=12.4$, $J=10.4$), 4.46 (d, 1H, $J=7.2$, H_1'), 4.38 (p, 1H, H_2), 4.12-4.02 (m, 2H, H_{1a} ,
20 H_2), 3.89 (pseudo t, 1H, $J=7.2$, H_{1b}), 3.79-3.6 (m, 5H), 3.50 (pseudo t, 2H), 1.46 (s, 3H), 1.40 (s, 3H); $^{13}\text{C-NMR}$ (CDCl_3) δ 138.529 (O), 138.370 (O), 138.066 (O), 138.013 (O), 128.432, 128.409, 128.129, 128.015, 127.901, 127.810, 127.734, 127.666, 109.399 (O),
25 103.824 (C_1'), 84-631 (C_3'), 82.120 (C_2'), 77.713 (C_4'), 75.748 (2), 75.058 (2), 74.891, 74.853, 74.315 (2), 73.495 (C_1), 70.317 (2), 68.762 (C_6'), 66.896 (C_3), 26.880 (3), 25.386 (3). Yield of the colorless, oily α -epimer (23 g, 0.035 mol) was 19%; $^1\text{H NMR}$ (CDCl_3)
30 δ 7.4-7.24 (m, 18H), 7.14 (m, 2H), 4.98 (d, 1H, $J=10.8$), 4.88-4.78 (m, 3H), 4.67 (d, 1H, $J=12$), 4.62 (d, 1H, $J=11.6$), 4.47 (d, 2H $J=11.6$), 4.37 (t, 1H, $J=6.4$), 4.07 (pseudo pentet, 1H), 3.96 (t, 1H, $J=8.8$), 3.8-3.54 (m, 9H), 1.43 (s, 3H), 1.37 (s, 3H); $^{13}\text{C NMR}$
35 (CDCl_3) δ 138.764 (O), 138.203 (O), 138.165 (O), 137.816 (O), 128.440, 128.387, 128.364, 128.030, 127.947,

127.916, 127.886, 127.696, 127.590, 109.422 (0),
97.482 (C₁'), 81.885 (1), 79.890 (1), 77.508 (1),
75.703 (2), 75.088 (2), 74.535 (1), 73.457 (2), 73.108
5 (2), 70.279 (1), 69.020 (2), 68.314 (2), 67.040 (2),
26.827 (3), 25.424 (3).

(2R) 1-O-(2,3,4,6-Tetra-O-benzyl- β -D-glucopyranosyl)-
glycerol 2. A 5-L three-necked morton flask fitted
10 with a mechanical stirrer, condenser, and stopper was
charged with compound 1 (50 g, 76.2 mmol) in 60%
aqueous acetic acid (2.5 L). The acidic solution was
refluxed for 1.5 hours at 103°C and then cooled to
room temperature. Distilled water (1.5 L) was added
15 to the solution causing precipitation of a white
solid. The acidic solution was extracted with
methylene chloride (4 x 1 L) which was subsequently
neutralized with sodium bicarbonate solution and
concentrated to a white solid. Trituration with
20 diethyl ether gave white product. The remaining
mother liquor was flash chromatographed (silica gel,
50% ethyl acetate/hexane) to give white solid product.
The combined yield (61.9 g, 0.101 mol) was 83 %, mp
101.5-102.4°C (lit 76-78°C); ¹H NMR (CDCl₃) δ .40-7.26
25 (m, 18H), 7.19 (t, J=3.5, 2H), 5.0-4.7 (m, 5H), 4.64-
4.5 (m, 3H), 4.46 (d, 1H, J=8.0, H₁'), 4.0-3.60 (m,
11H, H₁'s, H₂, H₃, H₃', H_{6a}', H_{6b}', H₄', H₅', H₂'), 2.55 (s,
2H, OH's); ¹³C NMR (CDCl₃) 38.529 (0), 138.332 (0),
137.952 (0), 137.740 (0), 128.531, 128.550, 128.478,
30 128.189, 128.114, 128.091, 127.931, 127.871,
127.749, 104.279 (C₁'), 84.654 (C₃'), 82.158 (C₂'),
77.819 (C₄'), 75.779 (2), 75.081 (2), 75.028 (2),
74.527 (C₅'), 73.571 (2), 72.207 (C₁'), 71.204 (C₂'),
68.883, (C₆'), 63.353 (C3).

35

(2S) [1-O-(2,3,4,6-Tetra-O-benzyl- β -D-glucopyranosyl)-3-O-tert-butyldimethylsilyl] glycerol 3. In a nitrogen-purged 100-mL round-bottomed flask fitted with a septum was dissolved diol 2 (9.0 g, 14.7 mmol), imidazole (2.05 g, 30.2 mmol), and t-butyl dimethylsilylchloride (2.28 g, 15.1 mmol) in anhyd DMF (45 mL). The reaction mixture was stirred under nitrogen for 2.5 days, transferred to a 1-L separatory funnel, and methylene chloride (250 mL) and water (250 mL) were added. The aqueous layer was extracted with methylene chloride (2 x 250 mL) and then the combined organic layers were washed with water (2 x 100 mL). After drying and concentration, purification by flash chromatography (silica gel, 33% ethyl acetate/hexane) gave a colorless oil (9.2 g, 12.6 mmol) in 88% yield; ^1H NMR (CDCl_3) δ 7.48-7.3 (m, 18H), 7.25-7.21 (m, 2H), 5.00 (d, 2H, $J=11.2$), 4.89 and 4.88 (overlapping doublets, 2H, $J=10.8$, $J=10.4$), 4.83 (d, 1H, $J=11.2$), 4.67 (d, 1H, $J=12.4$), 4.60 and 4.59 (overlapping doublets, 2H, $J=12.4$, $J=10.8$), 4.50 (d, 1H, $J=7.6$ Hz), 4.06-3.92 (m, 2H), 3.9-3.62 (m, 7H), 3.6-3.52 (m, 2H), 3.04 (s, 1H, OH), 0.978 (s, 9H), 0.142 (s, 6H); ^{13}C NMR (CDCl_3) δ 188.552 (O), 138.385 (O), 138.005 (O), 137.960 (O), 128.455, 128.440, 128.121, 128.060, 127.931, 127.863, 127.772, 127.734, 127.696, 104.377 (C1'), 84.692 (C3'), 82.219 (C2'), 77.804 (C4'), 75.771 (2), 75.073 (2), 74.959 (2), 74.717 (C5'), 73.541 (2), 73.078 (2), 71.060 (C2), 68.785 (C6'), 63.998 (C3), 25.970 (3), 18.668 (O), -5.299 (3).

(2S) [1-O-(2,3,4,6-Tetra-O-benzyl- β -D-glucopyranosyl)-2-O-palmitoyl-3-O-t-butyldimethylsilyl] glycerol 4. A nitrogen purged 500-mL round-bottomed flask fitted with a septum was charged with compound 3 (9.3 g, 12.8 mmol) and palmitic anhydride (6.94 g, 14.0 mmol) in

dry THF (200 mL). Dimethylaminopyridine (316 mg, 2.6 mmol) and triethylamine (2.04 mL, 14.7 mmol) were added, and the reaction was stirred under nitrogen for 12 h. The mixture was then transferred to a 2-L separatory funnel, and diethyl ether (500 mL) and water (500 mL) were added. The aqueous layer was filtered through Whatman No. 1 paper and extracted with diethyl ether (2 x 500 mL). After drying over magnesium sulfate, the combined organic layers were concentrated and purified by flash chromatography (silica gel, 14% ethyl acetate/hexane) to give a light yellow oil (12.1 g, 12.5 mmol) in 97% yield; ¹H NMR (CDCl₃) δ 7.40 (br. s, 20H), 5.15 (s, 1H), 4.98 (t, 2H), 4.84 (t, 2H), 4.76 (d, 1H), 4.67 (d, 1H), 4.59 (dd, 2H), 4.52 (d, 1H), 4.13 (dd, 1H), 3.84 (m, 6H), 3.67 (dd, 2H), 3.49 (t, 2H), 2.32 (t, 2H), 1.61 (m, 2H), 1.25 (br. s, 24H), 0.98 (s, 9H), 0.97 (s, 3H), 0.14 (s, 6H). ¹³C NMR (CDCl₃) δ 73.280, 138.597, 138.438, 138.127, 138.096, 128.379, 128.356, 128.333, 128.083, 127.977, 127.863, 127.780, 127.605, 127.582, 103.831, 84.556, 81.984, 77.721, 75.695, 75.020, 74.876, 74.603, 73.488, 72.904, 68.754, 67.821, 61.661, 34.428, 33.950, 31.941, 29.717, 29.687, 29.649, 29.619, 29.497, 29.459, 29.384, 29.300, 29.148, 25.826, 24.953, 22.716, 18.268, 14.159, -5.375.

(2R) [1-O-(2,3,4,6-Tetra-O-benzyl-β-D-glucopyranosyl)-2-Opalmitoyl] glycerol 5.

Procedure A. Compound 4 (34.0 g, 35.1 mmol) was dissolved in THF (1.4 L) in a 3-L three-necked Norton flask fitted with a mechanical stirrer, thermometer, and a 500-mL addition funnel. The solution was chilled to 0°C, and a solution of tetrabutylammonium fluoride (TBAF) (520 mL, 1.0 M in THF) which was buffered to pH=6.5 with acetic acid was added dropwise

through the addition funnel. The reaction mixture was stirred for 11 h at 0°C, left to sit at -15°C for 12 h, and stirred again for 4 h at rt. Water (100 mL) was added, and the solution was concentrated to 200 mL of solution. The concentrate was redissolved in methylene chloride (750 mL) in a 3-L separatory funnel and washed with water three times (750 mL, 2 x 500 mL). The combined aqueous layers were extracted with diethyl ether (500 mL). The combined organic layers were concentrated to give a red oil which was purified by flash chromatography (silica gel, 33-40% gradient of ethyl acetate/hexane). A white solid (28.0 g, 32.8 mmol) was obtained in 93% yield. ¹H NMR (CDCl₃) δ 7.36 (br. s, 20H), 5.06 (t, 1H), 4.96 (dd, 2H), 4.84 (dd, 2H), 4.75 (d, 1H), 4.59 (m, 2H), 4.53 (dd, 1H), 4.45 (dd, 1H), 4.14 (m, 2H), 3.91 (m, 2H), 3.78 (m, 6H), 2.80 (s, 1H), 1.64 (m, 2H), 1.27 (br. s, 26H), 0.90 (t, 3H).

Procedure B. Compound 4 (500 mg, 0.52 mmol) was dissolved in THF (20 mL) in a 100-mL three-necked round-bottomed flask fitted with two stoppers and a septum. Glacial acetic acid (9.5 mL) was added, and the solution was chilled to 0°C. A solution of TBAF (5.16 mL, 1.0 M in THF) was syringed into the chilled solution, and stirring was continued at 0°C for 8 h and then at rt for 25 hours. Methylene chloride (50 mL) was added, and the entire solution was transferred to a 250-mL separatory funnel where it was neutralized with 1M disodium phosphate solution (2 x 75 mL). The combined organic layers were rotary evaporated under reduced pressure and the concentrate was purified by flash chromatography (silica gel, 25-40% gradient of ethyl acetate/hexane), yielding a colorless oil (424 mg, 0.497 mmol, 95%) which later solidified upon standing; ¹H NMR (CDCl₃) δ 7.36 (br. s, 20H), 5.06 (t,

1H), 4.96 (dd, 2H), 4.84 (dd, 2H), 4.75 (d, 1H), 4.59 (m, 2H), 4.53 (dd, 1H), 4.45 (dd, 1H), 4.14 (m, 2H), 3.91 (m, 2H), 3.78 (m, 6H), 2.80 (s, 1H), 1.64 (m, 2H), 1.27 (br. s, 26H), 0.90 (t, 3H).

(28) [1-O-(2,3,4,6-Tetra-O-benzyl- β -D-glucopyranosyl)-3-O-palmitoyl] glycerol 6. Compound 4 (3.0 g, 3.1 mmol) was dissolved in THF (120 mL) in a 250-mL three-necked round-bottomed flask fitted with a 60-mL addition funnel, glass stopper, and septum. TBAF (54 mL, 1.0 M in THF) was added through the addition funnel over a 15 minute period. Glacial acetic acid (18 mL) measured in a graduated cylinder was then poured into the reaction mixture, and the solution was stirred for 45 minutes. The solution was concentrated under reduced pressure to approximately 30 mL of liquid and then redissolved in methylene chloride (150 mL). The organic layer was washed with water (3 x 120 mL) and neutralized with sodium bicarbonate solution (2 x 150 mL). The combined aqueous layers were extracted with methylene chloride (100 mL). The combined organic layers were dried over magnesium sulfate, filtered, and concentrated. The resulting dark red concentrate was purified by flash chromatography (silica gel, 25% ethyl acetate/hexane) to give 6 a colorless oil which corresponded to an upper TLC spot (1.3 g, 1.52 mmol) in 46% yield. ¹H NMR (CDCl₃) δ 7.36 (br. s, 20H), 4.95 (m, 2H), 4.86 (m, 3H), 4.64 (d, 1H), 4.58 (m, 2H), 4.47 (d, 1H), 4.16 (m, 1H), 3.96 (dd, 1H), 3.68 (m, 8H), 2.38 (t, 2H), 1.62 (m, 2H), 1.27 (br. s, 24H), 0.96 (t, 3H). Isolation of a lower TLC spot gave a white solid (400 mg, 0.469 mmol) in 15% yield which corresponded to compound 5; ¹H NMR (CDCl₃) δ 7.36 (br. s, 20H), 5.06 (t, 1H), 4.96 (dd, 2H), 4.84 (dd, 2H), 4.75 (d, 1H), 4.59 (m, 2H),

4.53 (dd, 1H), 4.45 (dd, 1H), 4.14 (m, 2H), 3.91 (m, 2H), 3.78 (m, 6H), 2.80(s, 1H), 1.64 (m, 2H), 1.27 (br. s, 26H), 0.90 (t, 3H).

5
Resilation of (2R) [1-O-(2,3,4,6-Tetra-O-benzyl- β -D
glucopyranosyl)-2-O-palmitoyl] glycerol 5. In a
nitrogen-purged 50-mL round-bottomed flask fitted with
a septum was placed compound 5 (318 mg, 0.373 mmol)
10 dissolved in DMF (8 mL). *tert*-Butyl-dimethylsilyl
chloride (281 mg, 1.86 mmol) and imidazole (254 mg,
3.73 mmol) were added, and the solution was stirred
for 22 h. Methylene chloride (50 mL) was added, and
the reaction mixture was transferred to a 250-mL
15 separatory funnel. The organic layer was washed with
water (50 ml), and then the aqueous layer was
extracted with methylene chloride (2 x 50 mL). The
pooled methylene chloride layers were washed with
water (2 x 75 mL), dried over magnesium sulfate, and
20 the filtered. The filtrate was concentrated and
purified by flash chromatography (silica gel, 14%
ethyl acetate/hexane) to give 4 as a yellow oil (239
mg, 0.247 mmol) in 66% yield.

25 Resilation of (2S) [1-O-(2,3,4,6-Tetra-O-benzyl- β -
-D-glucopyranosyl)-3-O-palmitoyl] glycerol 6. In a
nitrogen-purged 25-mL round-bottomed flask fitted with
a septum was placed compound 6 (176 mg, 0.206 mmol)
dissolved in anhyd DMF (5 mL). *tert*-
30 Butyldimethylsilyl chloride (155 mg, 1.03 mmol) and
imidazole (140 mg, 2.06 mmol) were added, and the
solution was stirred for 43 h. Methylene chloride (50
mL) was added, and the reaction mixture was
transferred to a 250-mL separatory funnel. The
35 organic layer was washed with water (50 mL). The
aqueous layer was extracted with methylene chloride (2

x 50 ml). The methylene chloride layers were pooled methylene and washed with water (2 x 75 mL), dried over magnesium sulfate, and then filtered. The
5 filtrate was concentrated and flash chromatographed (silica gel, 14% ethyl acetate/hexane) to give 7 as a light yellow oil (190 mg, 0.223 mmol) in 95% yield; ¹H NMR (CDCl₃) δ7.38 (br. s, 20H), 4.99 (dd, 2H), 4.85 (t, 2H), 4.78 (d, 1H), 4.68 (d, 1H), 4.61 (dd, 2H), 4.48
10 (d, 1H), 4.37 (d, 1H), 4.13 (s, 2H), 3.98 (m, 1H), 3.77 (m, 2H), 3.67 (m, 3H), 3.52 (m, 2H) 2.35 (t, 2H), 1.67 (m, 2H), 1.31 (br. s, 24H), 0.93 (s, 12H), 0.14 (s, 6H).

15 (2S) [1-O-(2,3,4,6-Tetra-O-benzyl-β-D-glucopyranosyl)-2-O-palmitoyl 3-O-(2-bromoethyl)benzylphosphoryl] glycerol 15.

Procedure A. In a nitrogen-purged 100-mL three-necked round-bottomed flask fitted with two stoppers and a
20 septum was dissolved freshly distilled 2-bromoethylphosphorodichloridate (1.72 g, 7.11 mmol) in diethyl ether (20 mL). The solution was chilled to 0°C, and triethylamine (8.15 mL, 58.5 mmol) was injected into the solution which caused precipitation
25 of a white solid. A solution of compound 5 (1.0 g, 1.17 mmol) in anhyd diethyl ether (55 ml) was injected into the chilled reaction mixture, and the ice bath was removed. The reaction was stirred for 30 minutes after which benzyl alcohol (1.21 mL, 11.7 mmol) was
30 injected into the reaction mixture. Stirring was continued at rt for 5 d. The reaction was then filtered through a fritted glass funnel, and the filtrate was concentrated. The orange concentrate was purified by flash chromatography (0-33% ethyl
35 acetate/hexane) to give 15 as a light yellow oil (566 mg, 0.501 mmol) in 43% yield; ¹H NMR (CDCl₃) δ7.38-7.25

(br. s, 23H), 7.16 (m, 2H), 5.26 (m, 1H), 5.10 (t, 2H), 4.94 (m, 2H), 4.81 (t, 3H), 4.71 (d, 1H), 4.61 (d, 1H), 4.55 (d, 2H), 4.39 (d, 1H), 4.25 (m, 4H),
5 4.08 (dd, 1H), 3.73 (m, 3H), 3.64 (dd, 2H), 3.42 (m, 4H), 2.27 (t, 2H), 1.58 (m, 2H), 1.25 (br. d, 24H), 0.89 (t, 3H).; ¹³C NMR (CDCl₃) δ173.210, 138.559, 138.362, 138.096, 138.074, 128.667, 128.622, 128.333, 128.318, 128.296, 127.962, 127.878, 127.757, 127.734,
10 127.696, 127.605, 127.522, 103.862, 84.540, 81.969, 71.652, 75.589, 74.937, 74.906, 74.686, 73.480, 70.469, 70.385, 69.680, 69.619, 68.777, 67.283, 66.099, 66.069, 34.170, 31.887, 29.657, 29.619, 29.596, 29.452, 29.315, 29.239, 29.080, 24.802,
15 22.647, 14.050.

Procedure B. In a nitrogen-purged 100-mL three-necked roundbottomed flask fitted with a thermometer, stopper, and septum was dissolved freshly distilled 2-bromoethylphosphorodichloridate (1.42 g, 5.85 mmol) in
20 methylene chloride (15 mL). The solution was chilled to 0°C, and compound 5 (1.0 g, 1.17 mmol) and a solution of N-methylmorpholine (1.28 mL, 11.7 mmol) dissolved in methylene chloride (35 mL) was injected into the solution over a 10 minute period. The
25 reaction mixture was stirred at 0°C for 5.5 h at which point a new TLC spot which co-spotted with secondary alcohol 6 appeared. Stirring was continued for another 30 minutes, and benzyl alcohol (1.21 ml, 11.7 mmol) was injected into the reaction. After 6 days of
30 stirring, the reaction mixture was transferred to a 500-mL separatory funnel, and methylene chloride (150 mL) and water (200 ml) were added. The layers were separated, and the organic layer was rotary evaporated under reduced pressure. The resulting oil was flash
35 chromatographed (silica gel, 33% ethyl acetate/hexane) to give 15 as a yellow oil (250 mg, 19%); ¹H NMR

(CDCl₃) δ 7.38 8-7.2 5 (br. s, 23H), 7.16 (m, 2H), 5.26 (m, 1H), 5.10 (t, 2H), 4.94 (m, 2H), 4.81 (t, 3H), 4.71 (d, 1H), 4.61 (d, 1H), 4.55 (d, 2H), 4.89 (d, 1H), 4.25 (m, 4H), 4.08 (dd, 1H), 3.73 (m, 3H), 3.64 (dd, 2H), 3.42 (m, 4H), 2.27 (t, 2H), 1.58 (m, 2H), 1.25 (br. d, 24H), 0.89 (t, 3H).; ¹³C NMR (CDCl₃) δ 173.210, 138.491, 138.286, 137.990, 137.975, 128.720, 128.652, 128.387, 128.364, 128.015, 127.954, 127.878, 127.810, 127.780, 127.727, 127.681, 127.613, 103.854, 84.495, 81.923, 77.781, 77.546, 75.688, 75.020, 74.808, 74.747, 73.473, 70.438, 69.642, 68.633, 67.322, 66.759, 66.129, 34.178, 31.925, 29.702, 29.664, 29.641, 29.490, 29.422, 29.368, 29.285, 29.103, 24.802, 22.700, 14.198.

(2S) [1-O-(2,3,4,6-Tetra-O-benzyl- β -D-glucopyranosyl)-2-O-palmitoyl-3-O-phosphatidylcholine] glycerol 16. A 45 mL Parr bomb equipped with a magnetic stirring bar was charged with a solution of phosphate 15 in toluene (10 mL). Condensed anhydrous trimethylamine (12 mL) was added quickly in one portion, and then the vessel was sealed and heated in an oil bath at 55°C for 24 h. The reaction mixture was concentrated to a viscous oil and triturated with ethyl ether, upon which a white precipitate formed. the precipitate was filtered off, washed with ether, and then the combined ethereal solutions were concentrated to a glassy solid. Purification of this residue using preparative TLC (2000 μ double elution with 75%,12.5%,12.5% methylene chloride/reethanol/ hexanes gave inner salt 16 as a glassy solid;

(2S) [β -D-glucopyranos-1-yl-2-O-palmitoyl-3-O-phosphatidylcholine] glycerol SP-19501. A solution of phosphatidylcholine 16 (200.4 mg, 0.197 mmol) in

reagent grade methanol (25 mL) was hydrogenated at 60 psi over 10% Pd/C (40 mg, 20 wt%). After 30 h, the catalyst was filtered off through celite and the methanol washings were combined and concentrated. The residue was dissolved in fresh methanol (25 mL) and resubjected to hydrogenation at 60 psi over 80 mg (40 wt%) of 10%Pd/C. After 48 h, the reaction was still incomplete. After filtration, washing of the catalyst, and concentration, the residue was subjected to hydrogenation using 400 mg (200 wt%) of Pd/C at 60 psi in methanol (25 mL). After 22h, the catalyst was filtered off through celite and the methanol filtrate and washings were combined and concentrated to afford 92.8 mg (71.6%) of (S) SP-19501 as a white solid; ¹H NMR (CD₃OD) δ 5.12 (br t, 0.5 H), 4.88 (br m, 4.5 H), 4.25 (br m, 2H), 4.12-3.57 (M, 12H), 3.4-3.1 (m containing singlet at 3.18, 12H), 2.3 (m, 2H), 1.55 (m, 2H), 1.24 (m, 22H), 0.86 (br t, 3H); ¹³C NMR (CD₃OD) δ 74.93, 104.80, 78.02, 77.93, 75.19, doublet at 71.53 and 71.49, doublet at 70.80 and 70.73, doublet at 67.79 and 67.74, multiplet at 67.50, 62.53, doublet at 60.56 and 60.52, triplet at 54.79, 34.88, 33.15, 30.85, 30.85, 30.66, 30.56, 30.46, 30.26, 26.10 and 26.03, 23.82, 14.56; ³¹P NMR (CD₃OD) δ 1.65.

(2S) 2,3-O-Isopropylidene-1-O-trifluoromethylsulfonyl-glycerol was prepared according to the method described for the corresponding (R) isomer in 92% yield and used immediately.

(2S) [1-O-(2,3,4,6-Tetra-O-benzyl-β-D-glucopyranosyl)-2,3-O-isopropylidene] glycerol 8. 2,3,4,6-Tetra-O-benzyl-D-glucopyranose (65 g, 0.12 mol) was dissolved in THF (800 mL) and chilled to -10°C in a nitrogen-purged 3-L three-necked morton flask fitted with a

- thermometer, stopper, and mechanical stirrer. Sodium hydride 60% in oil (33 g, 0.825 mol) was added in 4 increments over 10 minutes, and the solution was stirred for 1h. (S) 2,3-O-Isopropylidene-1-O-trifluoromethylsulfonylglycerol (0.15 mol) dissolved in THF (200 mL) was then dropped via an addition funnel into the reaction mixture over a 20 minute period at -10 to -15°C. The solution was stirred at -10 to -15°C for 6 hours. The reaction mixture was filtered through a short plug of silica gel and concentrated to an orange brown oil, 114 g. Purification of the crude by flash chromatography using 50% ethyl ether/hexanes gave 39.8 g (67.6%) of β epimer 8 as a white solid, along with 4 g (5.1 %) of a mixture of α and β epimers; mp of β anomer 85.7-87.2°C; ^1H NMR of β epimer (CDCl_3) δ 7.4-7.2 (m, 18H), 7.19-7.14 (m, 2H), 4.98-4.92 [overlapping doublets at 4.97 (J=10.8) and 4.94 (J=10.8), 2H], 4.82 (t, 2H, J=10.8), 4.73 (d, 1H, J=10.4), 4.63 (d, 1H, J=12.4), 4.58-4.51 [overlapping doublets at 4.55 (J=12) and 4.53 (J=10.8), 2H], 4.45 (dy 1H, J=7.2), 4.36 (p, 1H, H2), 4.08 (pseudo triplet, 1H), 3.94-3.89 [overlapping doublets at 3.92 (J=10) and 3.91 (J=9.6), 1H], 3.82-3.57 (m, 6H), 3.47 (pseudo triplet, 2H), 1.44 (s, 3H), 1.38 (s, 3H); ^{13}C NMR (CDCl_3) δ 138.569 (O), 138.384 (O), 138.006 (O), 138.021 (O), 128.341, 128.258, 127.962, 127.856, 127.765, 127.696, 127.620, 127.605, 109.467 (O), 103.869 (C1'), 84.586 (C3'), 82.075 (C2'), 77.705 (C4'), 75.680 (2), 75.005 (2), 74.815 (2 carbons, C2, C5'), 74.512 (1), 73.457 (C1), 71.151 (2), 68.785 (C6'), 67.017 (C3), 26.895 (3), 25.393 (3).
- (2S) 1-O-(2,3,4,6-Tetra-O-benzyl- β -D-glucopyranosyl)-glycerol 9. A suspension of 8 (20 g, 30.5 mmol) in

60% acetic acid (800) was heated to reflux for 1h. Workup was similar to that described for the (R) diol 2, providing 18 g (96% yield) of 9 as a white solid, which was of sufficient purity after trituration with ether for the subsequent step. Diol 9 could be recrystallized from ether/hexane, mp 89.6-90.90C; ¹H NMR (CDCl₃) δ7.38-7.27 (m, 18H), 7.16 (t, J=3.5, 2H), 4.98-4.74 (m, 5H), 4.61-4.5 (m, 3H), 4.42 (d, 1H, J=8.0, H₁'), 3.89-3.80 (m, 3H, H₁'s, H₂), 3.72-3.63 (m, 4H, H₃ H₃', H_{6b}'), 3.62-3.44 (m, 4H, H_{6a}', H₄', H₅', H₂'), 2.59 (s, 2H, OH's); ¹³C NMR (CDCl₃) δ138.370 (O), 138.119 (O), 137.78 (O), 137.69 (O), 128.462, 128.447, 128.432, 128.060, 128.038, 127.962, 127.894, 127.848, 127.810, 127.704, 104.195 (C₁'), 84.616 (C₃'), 82.037 (C₂'), 77.736 (C₄'), 75.733 (2), 75.043 (2, 2 carbons), 74.466 (C₅'), 73.480 (2), 72.312 (C₁), 70.772 (C₂), 68.731, (C₆'), 63.355 (C₃).

(2R) [1-O-(2,3,4,6-Tetra-O-benzyl-β-D-glucopyranosyl)-3-O-tert-butyldimethylsilyl] glycerol 10. In a nitrogen-purged 100-mL round-bottomed flask fitted with a septum was dissolved diol 9 (28.0 g, 45 mmol), imidazole (5.71 g, 90 mmol), and t-butyl dimethylsilylchloride (6.92 g, 45.3 mmol) in anhyd DMF (75 mL). The reaction mixture was stirred under nitrogen overnight, transferred to a 1-L separatory funnel, and chloroform (300 mL) and water (300 mL) were added. The aqueous layer was extracted with chloroform (2 x 100 mL) and then the combined organic layers were washed with water (3 x 100 mL). After drying (Na₂SO₄) and concentration, purification by flash chromatography (silica gel, 50% ethyl ether/hexanes) gave 10 as a colorless oil (29.5 g) in 90% yield;

(2R) [1-O-(2,3,4,6-Tetra-O-benzyl- β -D-glucopyranosyl)-2-Opalmitoyl-3-O-t-butyldimethylsilyl] glycerol 11. A mixture of 10 (22.2 g, 30.4 mmol), palmitic anhydride (16.5 g, 33.4 mmol), dimethylaminopyridine (741 mg, 6.08 mmol), triethylamine (3.78 g, 5.2 mL, 37.3 mmol) and anhyd THF (250 mL) was stirred under nitrogen at rt overnight. The mixture was poured into a 2-L separatory funnel, diluted with diethyl ether (500 mL) and water (500 mL), and the layers separated. The aqueous layer was filtered through Whatman No. 1 paper and extracted with more diethyl ether (2 x 500 mL). The combined ether layer was washed with water (3 x 200 mL) and then dried (MgSO_4). Following filtration, purification by flash chromatography (silica gel, 33% ethyl ether/hexane) gave 11 as a light yellow oil, 28.2 g, 96% yield);

Compound 11 could be carried on to the next transformation without chromatographic purification.

(2S) [1-O-(2,3,4,6-Tetra-O-benzyl- β -D-glucopyranosyl)-2-Opalmitoyl] glycerol 12. Crude 11 (30.4 mmol based on 10) was dissolved in THF (100 mL) and the solution was chilled to 0°C. A premixed solution of TBAF (520 mL, 1.0 M in THF) which was buffered to pH=6.37 with acetic acid was added dropwise via an addition funnel at 0°C for 1 h, and then at -15°C overnight. The reaction mixture was concentrated, water (100 mL) was added, and the resulting mixture was extracted with chloroform (3 x 300 mL). The combined chloroform layer was washed with water (4 x 500 mL), and then the combined aqueous layer was backextracted with diethyl ether (500 mL). After drying the combined organic layer over Na_2SO_4 , concentration gave a red oil which was purified by flash chromatography (50% ethyl ether/hexane). Evaporation of the product containing

fractions afforded 12 as a white solid (24.6 g, 94.6% yield for two steps);

- 5 (2R) [1-O-(2,3,4,6-Tetra-O-benzyl- β -D-glucopyranosyl)-3-Opalmitoyl] glycerol 13. Compound 11 (2.33 g, 2.41 mmol) was dissolved in THF (150 mL) in a 250-mL three-necked round-bottomed flask fitted with a 60-mL addition funnel, glass stopper, and septum. After
10 cooling the solution to 0°C, TBAF (24.1 mL, 1.0 M in THF) was added through the addition funnel over a 5 minute period. Glacial acetic acid (13.8 mL 241 mmol) was then poured into the reaction mixture to quench the reaction, and the resulting solution was stirred
15 for approximately 30 minutes. The reaction mixture was poured into a separatory funnel containing ice water (500 mL) and methylene chloride (200 mL). The layers were separated, and aqueous layer was extracted twice more with methylene chloride (100 mL portions)
20 and then the combined organic layer was washed with brine (400 mL). Following drying (MgSO_4), filtration, and then concentration, purification by flash chromatography using 1/5 EtOAc/ hexanes gave secondary alcohol 13, 0.96 g (46.8%), as a colorless oil;
25 Further elution gave 238 mg (11.6%) of primary alcohol 12; Also isolated was a mixture of the two alcohols in 5.3% yield.

- Resilylation of (2S) [1-O-(2,3,4,6-Tetra-O-benzyl- β -D glucopyranosyl)-2-O-palmitoyl] glycerol 12. The
30 identity of 12 was established by resilylation of 12 according to the procedure described above for the (R) isomer, compound 5.

35

- (2R) [1-O-(2,3,4,6-Tetra-O-benzyl- β -D-glucopyranosyl)-2-O-palmitoyl-3-O-(2-bromoethyl)benzylphosphoryl] glycerol 17. In a nitrogen-purged 1-L three-necked
5 morton flask fitted with two stoppers and a septum was dissolved freshly distilled 2-bromooethylphosphorodichloridate (17.2 g, 71.1 mmol) in anhyd diethyl ether (500 mL). The solution was chilled to 0°C and triethylamine (81.5 mL, 0.585 mol)
10 was injected into the solution, causing precipitation of a white solid. A solution of 12 (10.0 g, 11.7 mmol) dissolved in diethyl ether (250 mL) was cannulated into the morton flask, and the solution was stirred for 1.5 hours. TLC showed disappearance of
15 12. Benzyl alcohol (12.1 mL, 0.117 mol) was injected into the reaction mixture, and stirring was continued at rt for 16 h. The reaction mixture was then filtered through a fritted glass funnel. Filtrate was then concentrated and purified by flash chromatography
20 twice. First chromatography (silica gel, 33% ethyl acetate/hexane) and second chromatography (silica gel, 25% ethyl acetate/hexane) gave 17 as a light oil (5.5 g) in 42% yield;
- 25 (2S) [1-O-(2,3,4,6-Tetra-O-benzyl- β -D-glucopyranosyl)-2-O-palmitoyl-3-O-phosphatidylcholine] glycerol 18. A 45 mL Parr bomb was equipped with a magnetic stir bar and then charged with a solution of 17 (1.17 g, 1.04 mmol) in benzene (15 mL).
30 Anhyd trimethylamine (15 mL, 0.145 mmol) which had been condensed at -78°C was quickly poured into the reaction vessel, and the bomb was sealed. The reaction was stirred at 55°C in an oil bath for 24 hours behind a blast shield. The bomb vessel was then
35 cooled to -78°C, opened, and left in a hood to evaporate trimethylamine. The remaining solution was

rotary evaporated under reduced pressure, and the oily concentrate was dissolved in methylene chloride and purified by preparative TLC (2000 μ). Double elution with 75%; 12.5%:12.5% methylene chloride/methanol/hexane gave inner salt 18 as an opaque glassy solid (223 mg, 21%). $^1\text{H-NMR}$ (CDCl_3) δ 6.32 (br. s, 20H), 5.21 (m, 1H), 4.90 (dd, 2H), 4.82 (m, 2H), 4.64 (m, 2H), 4.50 (t, 2H), 4.42 (d, 1H), 4.22 (br. s, 3H), 3.95 (s, 2H), 3.72 (s, 2H), 3.62 (t, 2H), 3.55 (s, 1H), 3.40 (m, 4H), 3.10 (s, 9H), 2.19 (m, 2H), 1.47 (m, 2H), 1.20 (br. d, 24H), 0.87 (t, 3H). $^{13}\text{C-NMR}$ (CDCl_3) δ 73.393, 138.453, 138.377, 138.051, 137.998, 128.470, 128.417, 128.356, 128.318, 128.235, 128.053, 128.007, 127.947, 127.856, 127.780, 127.719, 127.636, 127.567, 103.899, 84.472, 81.984, 77.478, 77.394, 77.311, 77.190, 75.672, 74.944, 74.550, 73.336, 68.663, 68.489, 59.158, 59.135, 54.409, 54.349, 34.246, 31.902, 29.710, 29.664, 29.535, 29.353, 29.330, 29.148, 24.7871 22.678, 14.121.

(2R) [β -D-glucopyranos-1-yl-2-O-palmitoyl-3-O-phosphatidylcholine] glycerol SP-19501. A solution of phosphatidylcholine 18 (130 mg, 0.127 mmol) in reagent grade methanol (25 mL) was hydrogenated at 60 psi over 10% Pd/C (52 mg, 40 wt%). After 23 h, TLC showed an incomplete reaction. The catalyst was filtered off through celite and the methanol washings were combined and concentrated. The residue was dissolved in fresh methanol (25 mL) and resubjected to hydrogenation at 60 psi over 240 mg (185 wt%) of 10% Pd/C. After 20 h, the reaction was complete by TLC. The catalyst was filtered off through celite and the methanol filtrate and washings were combined and concentrated to afford 64.0 mg (76.6%) of (R) SP-19501 as a white solid; $^1\text{H-NMR}$ (CDCl_3) δ 5.19 (m, 1H), 4.97 (s, OH + HDO), 4.34-

4.26 (br m, 2H), 4.16-3.95 (m, 3H), 3.9-3.6 (m, 6H),
3.42-3.14 (multiplet containing singlet at 3.24, 13H),
2.37 (t, J=7.6, 2H), 1.62 (pseudo t, 2H), 1.31 (m,
5 24H), 0.92 (t, J=7.2, 3H); ¹³C NMR (CD₃OD) δ 175.02,
104.88, 78.07, 78.04, 75.04, 72.97, 72.89, 71.52,
68.56, multiplet at 67.50, doublet at 64.99 and 64.94,
62.65, doublet at 60.52 and 60.48, triplet at 54.74
(J=3.1), 35.14, 33.13, 30.85, 30.69, 30.54, 30.29,
10 26.01, 23.79, 14.50; ³¹P NMR (CD₃OD) δ 1.35

(2R)1-[Benzyl-(2'-bromoethyl)-phosphpropyl]-2,3-
isopropylidene glycerol (19).

2-Bromoethylphosphodichloridate (20.0 g, 0.08 mol) was
15 dissolved in CC14 (50 ml) in a nitrogen-purged 0.5 L
three-necked flask fitted with a magnetic stir bar,
thermometer, and a 125-ml addition funnel. The
solution was chilled to 0°C, and to this stirred
solution was added dropwise the solution of (S)-form
20 solketal (10.7 g, 98 mol %) and N-methyl-morpholine
(8.22 g, 98 mol %) in CCL4 (25 ml). After 2 hours TLC
showed disappearance of solketal. To the reaction
mixture was added dropwise the solution of benzyl
alcohol (44.6 g, 500 mol %) and N-methylmorpholine
25 (8.38 g, 100 mol %). The reaction mixture was stirred
under nitrogen for 60 hours at room temperature. TLC
showed the complete reaction. The reaction mixture
was filtered through Shott filter #C, and the solution
was rotary evaporated to volume near 70 ml and
30 purified by flash chromatography (silica gel, diethyl
ether) to give colorless oil (15.1 g, 0.04 mol) in 45
% yield; ¹H NMR (CDCl₃) δ ppm: 7.40 (br. s 5 H), 5.2 (d,
2 H), 4.3 (br.s, 3 H), 4.0 (br.s, 3 H), 3.85 (br.s, 1
H), 3.2 (s, 2 H), 1.4 (d, 6H); ¹³C NMR (CDCl₃):
35 128.743, 128.682, 128.645, 128.114, 128.076, 109.885,
77.364, 77.046, 76.727, 73.920, 73.837, 69.771,

69.710, 67.760, 67.707, 67.654, 66.774, 66.721,
65.955, 29.353, 29.277, 26.683, 25.204; ³¹P NMR
(CDCL₃): -1.108.

5
(2R)1-[Benzyl-(2'-bromoethyl)-phosphoroyl] 1-2,3-
dihydroxy glycerol (20). A nitrogen purged 1 L round-
bottomed flask fitted with septum was charged with
compound 19 (19.5 g, 0.048 mol) in dry THF (50 ml) and
10 the solution of 1 M H₃PO₄ (800ml) was added. The
reaction mixture was stirred under nitrogen by room
temperature for 15 hours. TLC showed the completeness
of the reaction. Then the reaction mixture was
transferred to a 2 L separatory funnel. The acidic
15 layer was extracted with ethyl acetate (7 x 450 ml).
The combined organic extract was washed with water (2
x 850 ml). After drying over sodium sulfate it was
rotary evaporated and dried in high vacuo for 10 hours
to give a colorless oil (14 g, 0.04 mol %) in 80 %
20 yield; ¹H NMR (CDCL₃), δppm: 7.38 (br.s, 5H), 5.2 (d,
2H), 4.25-3.8 (multiplet, 6 H), 3.7-3.25 (br.m 5 H) ¹³C
NMR (CDCL₃): 77.789, 77.774, 77.349, 77.030, 76.712,
70.522, 70.491, 70.461, 70.431, 70.097, 70.044,
68.898, 68.883, 68.822,
25 67.085, 67.032, 62.617, 62.496, 42.363, 42.280; ³¹P NMR
(CDCL₃): -0.485 (85 % H₃PO₄).

(2R)1-[Benzyl-(2'-bromoethyl)-phosphoroyl-2-hydroxy-3-
O-triphenylmethyl glycerol (21). To a stirred
30 solution of diol 20 (8.0 g, 21.6 mmol) in DMF (16 ml)
was added diisopropylethylamine (4 ml, 105 mol %)
followed by addition of trityl chloride (6.4 g, 105
mol %). After 40 hours at room temperature under
nitrogen the reaction was complete by TLC. The
35 reaction mixture was diluted twice with water and
extracted with diethyl ether (4 x 100 ml). The

combined extract was dried over sodium sulfate, concentrated and purified by flash chromatography silica gel, ethyl acetate:hexane,1:1) to give 21 as a
5 light oil 6.9 g (52.1 %); ¹H NMR (CDCL₃) δ.2-7.5 (br.m, 20 h), 5.07 (t, 2H), 4.12-4.26 (m,4H), 3.44 (dd 2H), 2.05 (s, 1H), 1.26 (t, 1H). ¹³C NMR (CDCL₃): 138.772, 124.017, 123.911, 123.820, 123.342, 123.266, 123-152, 122.462, 122.417, 72.593, 72.274, 71.955, 65.105,
10 65.090, 65.030, 65.014, 64.954, 64.893, 62.109, 62.056, 58.877, 24.680, 24.604. ³¹P NMR (CDCL₃) -0.158 (s).

(2R) 1-[Benzyl-(2'-bromoethyl)-phosphoroyl-2-O-
15 palmitoyl-3-O-triphenyl methyl glycerol (22). To a stirred solution of compound 21 (6.9 g, 11.3 mmol) in dry THF (90 ml) was added triethylamine (1.79 ml, 1 10,mol %), palmitic anhydride (6.13 g, 1 10 mol%) and dimethylaminopyridine (276 mg, 20 mol%). The reaction
20 was stirred under nitrogen for 3 hours untill TLC showed disappearance of the starting material 21. The reaction mixture was rotary evaporated to a small volume and purified by flash chromatography (silica gel, diethyl ether:hexane, 1:3 to elute UV-nonactive
25 impurities, diethyl ether:hexane,1:1 to elute compound 22). Yield 8.8 g (92.6%, colorless oil, ¹H NMR (CDCL₃) 7.41-7.22 (m, 20 H), 5.20 (d, 1H), 5.04 (t, 2H), 4.23 (m, 4H), 3.58 (s, 1H), 3.41 (s, 1H), 3.23 (s, 2H), 2.33 (t, 2H), 1.62 (m, 3H), 1.24 (s, 24H), 0.88 (t,
30 3H). ¹³C NMR 172.984, 143.407, 143.285, 128.675, 128.607, 128.576, 128.523, 127.985, 127.848, 127.180, 127.135, 86.672, 77.319, 77.000, 76.681, 70.901, 70.818, 69.604, 69.581, 66.463, 66.440, 61.828, 34.284, 31.894, 29.672, 29.634, 29.611, 29.437,
35 29.346, 29.285, 29.247, 29.232, 29.141, 29.095, 24.832, 22.678, 14.121. ³²P NMR (CDCL₃)-1.327.

(2R)1-[Benzyl(2'- bromoethyl)-phosphoro]-yl-2-O-palmitoyl-3-hydroxy glycerol 23. Procedure A To a stirred solution of compound 22 (3.2 g, 3.76 mmol) in 45 ml THF was added 45 ml 96% formic acid. After 2 hours at room temperature the reaction was complete by TLC. The reaction mixture was diluted twice with water, neutralized with sodium bicarbonate (3 x 300 ml). The combined extract was washed with water, dried over sodium sulfate, rotary evaporated to a small volume and purified by flash chromatography (silica gel, ethyl acetate: hexane, 1:3 to elute less polar impurities, ethyl acetate:hexane, 1:1 to elute compound 23. Yield 1.65 g (72.7 %), colorless oil.

Procedure B. A nitrogen purged 0.5 L round-bottomed flask fitted with condenser was charged with compound 22 (1 g, 1.17 mmol) in dry benzene (230 ml) in the presence of anhydrous CuSO₄ (17.6 g). The reaction mixture was stirred at room temperature for 15 hours and then reflux for 2 hours until the reaction was complete by TLC. The CuSO₄ was filtered off through Shott filter #C and concentrated in vacuo and purified by flash chromatography (silica gel, ethyl acetate/hexane, 1:1) to give a light yellow oil (0.47 g, 0.77 mmol) in 66 % yield. ¹H NMR (CDCl₃): α 7.40 (br. s, 5H), 5.2 (d, 2H), 4.2 (mult, 8H), 2.32 (t, 1H), 1.62 (pseudo t, 2H), 1.31 (m, 24H), 0.88 (t, 3H). ¹³C NMR (CDCl₃): 130.898, 128.872, 128.789, 128.698, 128.538, 128.516, 128.114, 127.886, 126.968, 77.326, 77.008, 76.689, 70.097, 70.074, 69.012, 68.951, 68.633, 68.604, 68.572, 68.542, 67.085, 67.047, 67.032, 66.994, 65.272, 64.195, 62.731, 62.716, 34.041, 31.902, 29.672, 29.588, 29.505, 29.444, 29.338, 29.239, 29.103, 24.749, 22.670, 14.113. ³¹P NMR: - 3.069 (85% H₃PO₄).

(2R)-1-O-(2,3,4,6-Tetra-O-benzyl- β -D-gluco-pyranosyl)-
2'-O-palmitoyl-3'-O-[benzyl(2''-bromoethyl)-
phosphoril]-glycerol (17). To a stirred solution of
5 O-(α -D-glucopyranosyl) trichloroacetimidate (24) (390
mg, 115 mol %) in dry methylene chloride (3 ml) was
added dropwise a solution of compound 6 (300 mg, 0.49
mmol) and boron trifluoride etherate (70 mg, 100 mol
%) in dry methylene chloride (3 ml). The reaction
10 mixture was stirred under nitrogen for 2 hours at room
temperature, then more compound 24 (100 mg, 35 mol %)
was added to bring the reaction to the end. After 4h,
the reaction mixture was evaporated to a small volume
and separated by flash chromatography (silica gel,
15 diethyl ether/hexane, 1:3) to give compound 17 as a
colorless oil (120 mg, 22%), which was identical to
the material described earlier.

6.4. Antifungal Activity

20 The antifungal activity of the isolated
phosphocholine fraction was determined *in vitro* by
using three fungal cultures --*Candida albicans*,
Cryptococcus neoformans and *Aspergillus fumigatus*.
The method used to determine *in vitro* antifungal
25 activity is discussed in McGinnis, M.R., *Laboratory
Handbook of Medical Mycology*, Academic Press, New
York, London, p661 (1980); and Droughet E., Dupont,
B., Improvisi, L., Vivian, M.A. and Tortorano, A.M.,
"Disc agar diffusion and microplate automatized
30 technics for *in vitro* evaluation of antifungal agents
on yeast and sporulated pathogenic fungi" in *In Vitro
and In Vivo Evaluation of Antifungal Agents*, Eds.
Iwata, K. and Vanden Bossche, H., Elsevier Science
Publishers, New York, Oxford p303 (1986).

35

The minimum inhibitory concentration (MIC) and the minimum fungicidal concentration (MFC) are summarized in the table 1 below.

5

<u>Fungus Culture</u>	<u>MIC (ug/ml)</u>	<u>MFC (ug/ml)</u>
C. albicans	0.8	
C. neoformans	< 0.1	
A. fumigatus	< 0.1	<0.4-0.8

10


These results clearly indicate the significant antifungal activity of the isolated fraction containing against a variety of fungal cultures.

15 6.5. Antifungal Activities of the Phosphocholine Derivatives Class

A series of related analogs to 2-palmitoyl-1-0-glucopyranosyllysolecithin obtained commercially from Avanti Biolipids have also been found to have high
20 antifungal activities. A summary of the antifungal screening test is shown in table 2. The analog compounds were tested for their activity against C. albicans, C. neoformans, A. fumigatus and T. rubrum. Partial inhibition of the fungus of between
25 25 to 75% was measured along with the total inhibition (MIC) by these analog compounds. A description of the partial inhibition measurement can be found in R. L. Stiller, et al *The Journal of Infectious Diseases*, 147, No. 6 (1983). The structure of these analog
30 compounds is as follows.

35

5



10

15

wherein R is the group identified in table 2.

20

25

30

35

Table 2		Test Results from Antifungal Screening Laboratory											
	Lecithin	MIC ($\mu\text{g/ml}$)						Partial inhibition ($\mu\text{g/ml}$)					
		CA	CN	AF	TR	CA	CN	AF	TR	CA	CN	AF	TR
5	L-a-Lysophosphatidylcholine Heptadecanoyl (C17:0)	63	16	>1000	63	16	n/a	31	31	16	n/a	31	31
10	L-a-Lysophosphatidylethanol Amine, Oleoyl (C18:1, [cis]-9)	>100	>100	>100	>100	>100	100	>100	>100	>100	100	>100	>100
15	L-a-Lysophosphatidylcholine Decanoyl (C:10)	>500	500	>500	>500	250	250	>500	>500	250	250	>500	>500
	L-a-Lysophosphatidylcholine Lauroyl (C12:0)	500	125	125	250	n/a	31	n/a	63	n/a	31	n/a	63
20	L-a-Lysophosphatidylcholine Myristoyl (C14:0)	31	31	125	31	n/a	n/a	31	n/a	n/a	n/a	31	n/a
	L-a-Lysophosphatidylcholine Stearoyl (C18:0)	>250	>250	>250	>250	>250	31	>250	>250	>250	31	>250	>250

L- α -Lysophosphatidylcholine Oleoyl (C18:1,[cis]-9)	31	31	63	31	n/a	n/a	31	n/a
L- α -Lysophosphatidylcholine Palmitoyl (C16:0)	31	31	63	500	n/a	63	31	63
L- α -Lysophosphatidyl inositol	>100	>100	100	>100	>100	100	n/a	>100

5

10

15

20

25

6.6. Toxicity

The toxicity of the isolated phosphocholine derivative fraction is low, based on tests with Hep 2 cells indicating an ID₅₀ of greater than 1000 ug/ml. The method used in determining cytotoxicity is discussed in Mosmann, T., "Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays", *J. Immun. Methods*, 65, 55-63, 1986.

The isolated fraction having the above-described *in vitro* antifungal activity and low toxicity is expected to similarly exhibit significant *in vivo* antifungal activity against fungal infections which are dermatophytic, systemic, ophthalmic and vaginal. Other human and animal infections treatable with the compounds of the present invention include aspergilliosis, candidiasis, and cryptococcus infections.

It is expected that the same isolated fraction would be useful in treating fungal infestation in plants as well.


It is apparent that many modifications and variations of this invention may be made without departing from the spirit and scope thereof. The specific embodiments described are given by way of example only and the invention is limited only by the terms of the appended claims.

A number of references are cited in the present specification, the entire disclosure of each of which is incorporated by reference herein, in its entirety.

CLAIMS

What is claimed is:

1. A phosphocholine derivative having the
5 structure of:



15


wherein one of R or R' is a sugar moiety,
and the other is an acyl or sugar moiety.

2. The phosphocholine derivative according to
20 claim 1 wherein the sugar moiety is selected from the
group consisting of glucose, galactose, arabinose,
mannose, rhamnose, and other naturally occurring
sugars.

- 25 3. The phosphocholine derivative according to
claim 1 having the structure of:


30

35



2-Palmitoyl-1-O-glucopyranosyllysolecithin

15 4. An antifungal composition capable of inhibiting fungal growth when administered to warm blooded animals or plants comprising an effective amount of a phosphocholine derivatives having the structure of:



30 wherein one of R or R' is a sugar moiety, and the other is an acyl or sugar moiety.

35 5. A composition comprising a phosphocholine derivative obtained from *Irlbachia alata* characterized by:

(a) IR spectrum having peaks at approximately 1060, 1220, 1475, 1600-1700, 2850, 2950, and 3400 cm^{-1} ;

5 (b) ^1H NMR spectrum having major peaks at δ 1.2, 1.4, 1.7, 3.1, 3.5, 3.7 and 4.3; and

(c) FAB/MB mass spectrum having major peaks (>40%) at m/z 657, 612, 587, 586, 555, 493, 491, 475, 403, 277, 233, 201, 194, 179, 168, 165 and 163.

10

6. A composition comprising a phosphocholine derivative according to claim 5, characterized by:

(a) IR spectrum substantially illustrated in Fig. 1;

15 (b) ^1H NMR spectrum substantially illustrated in Fig. 2; and

(c) FAB/MB mass spectrum substantially illustrated in Fig. 3.

(d) HRMS (FAB^+) spectrum having a molecular
20 ion at 673.4669 amu.

7. A composition comprising a phosphocholine derivative obtained from *Irlbachia alata* by a method which comprises:


25 (a) extracting the whole plant, the leaves, the stems, the roots or the latex of the plant *Irlbachia alata* with a lower alcohol of about 1-3 carbons, acetone, water or other water miscible solvent
30 or combinations thereof to obtain an aqueous soluble fraction;

(b) subjecting the aqueous fraction to butanol extraction and the butanol-soluble fraction to gel filtration
35 using water and/or water and a water miscible solvent with or without a

5 buffer as the mobile phase; or to
reversed phase column chromatography
using water, and/or water and a water
miscible solvent as the mobile-phase;
or to gel permeation chromatography
using water and/or water and water
miscible solvent and acetonitrile with
or without a buffer as the mobile
10 phase; or combination thereof and
(c) collecting the fractions detected by
NMR spectroscopy.


8. A pharmaceutical composition which is useful
15 in treating a fungal infection when administered to a
warm-blooded animal a therapeutically effective amount
of an antifungal agent comprising a phosphocholine
derivative having the structure of:

20




30 wherein one of R or R' is a sugar moiety,
and the other is an acyl or sugar moiety.

9. A pharmaceutical composition which is useful
in treating a fungal infestation when administered to
a plant comprising an effective amount of an
35 antifungal agent comprising a phosphocholine
derivative having the structure of:



wherein one of R or R' is a sugar moiety,
 15 and the other is an acyl moiety or sugar moiety.

10. A pharmaceutical composition which is useful
 in treating a fungal infection when administered to a
 warm-blooded animal comprising a therapeutically
 20 effective amount of an antifungal agent comprising a
 compound having the structure of:



30 where Q is C2 to C30 alkyl, alkenyl, alkynyl, branched
 alkyl, branched alkenyl, or branched alkynyl;

Z is oxygen or sulfur; X and Y are independent
 oxygen, sulfur, CH₂, CF₂, or N-R₁;

A, B, and T are independently alkyl, alkenyl,
 35 alkynyl, branched alkyl, branched alkenyl, or branched
 alkynyl radicals of C1 to C20 chain lengths; are
 independently or together cycloalkyl or bridged

cycloalkyl radicals of ring size C3 to C20, or
cylcoalkenyl, bridged cycloalkenyl or
cyclo(polyene)radicals of ring size C4 to C20,
5 cycloalkynyl, bridged cycloalkenyl or
cyclo(polyalkynyl)radicals of ring size C8 to C20;


D is oxygen, sulfur, CH₂, CF₂, or N-R₂;

F is alkyl, alkenyl, alkynyl, branched alkyl,
branched alkenyl, branched alkynyl, cycloalkyl, bridged
10 cycloalkyl, cycloalkenyl or cycloalkynyl radicals
containing C1 to C20 carbon atoms;

R₁ and R₂ are independently hydrogen, alkyl,
alkenyl, alkynyl, branched alkyl, branched alkenyl,
branched alkynyl, cycloalkyl, bridged cycloalkyl,
15 cycloalkenyl, bridged cycloalkenyl, or cycloalkynyl
radicals containing C1 to C20 carbon atoms, or a
protecting group.

11. A pharmaceutical composition which is useful
20 in treating a fungal infection when administered to a
warm-blooded animal comprising a therapeutically
effective amount of an antifungal agent comprising a
compound having the structure of:

25



30

where AA, BB, DD are independent of each other or
equal to each other, the central carbon atom can be
either the R and S optical stereoisomer or a mixture
of R and S stereoisomers, and where AA, BB, and CC are
as follows:


35

AA is A-J with A attached to the carbon atom of
the three carbon central unit and J is defined below;

BB is B-Y, with B attached to the carbon atom of the three carbon central unit and Y is hydrogen, alkyl, alkenyl, alkynyl, poly(alkenyl), poly(alkynyl),
 5 or poly(alkenoalkynyl) radicals comprised of C1 to C20 carbon atoms; chain lengths or alkanoyl, alkenoyl, alkynoyl, poly(alken)oyl, poly(alkyn)oyl, poly(alkenoalkyn)oyl radicals of C2 to C20 chain lengths, alkyloxy, alkenyloxy, alkynyloxy,
 10 poly(alkenyl)oxy, or poly(alkynyl)oxy, poly(alkenoalkynyl)oxy radicals comprised of C1 to C20 carbon atoms;

DD is

15



; wherein

20


A is oxygen, sulfur, CH₂, CF₂ or N-R₁;

B is oxygen, sulfur, CH₂, CF₂ or N-R₂;

D is oxygen, sulfur, CH₂, CF₂ or N-R₃;

J is a furanose or pyranose radical of the type:

25






30

where X is oxygen, sulfur, CH₂, CF₂ or N-R₄;

F, K, L and M are independently hydrogen, hydroxyl, a protected hydroxyl, alkyloxy, thiol, alkylthio, arylthio, alkylsulfonyl, arylsulfonyl,
 35 amino, ammonium, alkylamino, alkylammonium, dialkylamino, dialkylammonium, trialkylamino,



- trialkylammonium where the alkyl chain on nitrogen is comprised of C1 to C20 carbon atoms; or alkyl, alkenyl, or alkynyl radicals comprised of C1 to C20
- 5 carbon atoms;
- Z is oxygen or sulfur;
- E is oxygen, sulfur, CH₂, CF₂, or N-R₅;
- G is alkyl, branched alkyl, cycloalkyl or bridged cycloalkyl radicals of C1 to C20 chain lengths;
- 10 Q is halogen, hydroxyl, protected hydroxyl utilizing a protecting group, O-arylsulfonyl-, O-alkylsulfonyl- or O-(perfluoroalkyl)sulfonyloxy, amino, ammonium, alkylamino, alkylammonium, dialkylamino, dialkylammonium, trialkylamino, or
- 15 trialkylammonium where the alkyl chains on nitrogen are C1 to C20; or Q=NR₁R₂R₃, where R₁, R₂, or R₃ can independently or together be a mixture of alkyl groups of C1 to C20 in chain length and a protecting group and R₁ can equal R₂, R₂ can equal R₃, or R₁ can equal R₃
- 20 which can equal R₃;
- R₁, R₂, R₃, R₄ and R₅ are independently alkyl, alkenyl, alkynyl, branched alkyl, branched alkenyl, branched alkynyl, cycloalkyl, bridged cycloalkyl, cycloalkenyl or cycloalkynyl radicals of C1 to C20
- 25 chain lengths, or any protecting group;
- and where W₁ and W₂ are P(-OR), (with R being phenyl, phenylmethyl, or negatively-charged oxygen), S=O, carbon, or sulfur, provided that if W₁ is not P(-OR) W₂ is P(-OR) and provided that if J is furanose
- 30 or pyranose radical then W₂ is P(-OR).

12. A pharmaceutical composition which is useful
in treating a fungal infection according to claim 11,
comprising a therapeutically effective amount of an
antifungal agent comprising a compound having the
structure of:



30


35




where R_1 is phenyl or phenylmethyl, hydrogen, or
 20 nil;
 R_2 is hydrogen phenylmethyl or any protecting
 group which can be cleaved by hydrogenolysis;
 Q is halogen, hydroxyl, protected hydroxyl
 utilizing a protecting group, O-arylsulfonyl-, O-
 25 alkylsulfonyl- or O-(perfluoroalkyl)sulfonyloxy,
 amino, ammonium, alkylamino, alkylammonium,
 dialkylamino, dialkylammonium, trialkylamino, or
 trialkylammonium where the alkyl chains on nitrogen
 are C1 to C20; or $Q = NR_1R_2R_3$, where R_1 , R_2 , or R_3 can
 30 independently or together be a mixture of alkyl groups
 of C1 to C20 in chain length and a protecting group
 and R_1 can equal R_2 , R_2 can equal R_3 , or R_1 can equal R_3
 which can equal R_3 .

35 13. A pharmaceutical composition which is useful
 in treating a fungal infection according to claim 11,


comprising a therapeutically effective amount of an antifungal agent comprising a compound having the structure of:





- 15 where R_1 is phenyl, phenylmethyl, hydrogen, or nil;
 R_2 is hydrogen, phenylmethyl or a protecting group cleavable by hydrogenolysis;
 R_3 is hydrogen or a protecting group;
 20 and Q is a halogen, hydroxyl, O-arylsulfonyl-, O-alkylsulfonyl- or O-(perfluoroalkyl)sulfonyloxy;

14. A pharmaceutical composition which is useful
 25 in treating a fungal infection according to claim 11, comprising a therapeutically effective amount of an antifungal agent comprising a compound having the structure of:




- 35 where R_1 is phenyl, phenylmethyl, hydrogen or nil;

R_2 is a protecting group, or hydrogen if R_1 is not hydrogen;

Q is a halogen, hydroxyl, O-arylsulfonyl-, O-alkylsulfonyl- or O-(perfluoroalkyl)sulfonyloxy.

15. A compound having the structure of:



15 where Q is C2 to C30 alkyl, alkenyl, alkynyl, branched alkyl, branched alkenyl, or branched alkynyl;

Z is oxygen or sulfure; X and Y are independently oxygen, sulfur, CH_2 , CF_2 , or N-R_1 ;

A , B , and T are independently alkyl, alkenyl, alkynyl, branched alkyl, branched alkenyl, or branched alkynyl radicals of C1 to C20 chain lengths; are independently or together cycloalkyl or bridged cycloalkyl radicals of ring size C3 to C20, or cycloalkenyl or cyclo(polyene)radicals of ring size C4 to C20, cycloalkynyl or cyclo(polyalkynyl)radicals of ring size C8 to C20;


D is oxygen, sulfur, CH_2 , CF_2 , or N-R_2 ;

F is alkyl, alkenyl, alkynyl, branched alkyl, branched alkenyl, branched alkynyl, cycloalkyl, bridged cycloalkyl, cycloalkenyl or cycloalkynyl radicals containing C1 to C20 carbon atoms;

R_1 and R_2 are independently hydrogen, alkyl, alkenyl, alkynyl, branched alkyl, branched alkenyl, branched alkynyl, cycloalkyl, bridged cycloalkyl, cycloalkenyl, bridged cycloalkenyl or cycloalkynyl

radicals containing C1 to C20 carbon atoms, or a protecting group.

16. A compound according to claim 6 having the structure of:



17. A method of synthesizing a phosphocholine derivative comprising the steps of:

(a) phosphorylating an alcohol with a halo-alkyl containing phosphorylating agent; and

(b) displacing the halide by an amine to produce the phosphocholine derivative.

18. A method of synthesizing a lysolecithin comprising the steps of:

(a) phosphorylating a glycosylating an acetenide derivative of glycerol;

(b) deprotecting the phosphorylated or glycosylated product; and

(c) alkylating or esterifying the deprotected product to form a lysolecithin.

1/3




FIG. 1

2/3




FIG. 2

SAMPLE		DATE, & UP	
date	Apr 7 03 dn		HE
delivered	Apr 0300 dof		-554.0
title	Photo/Mar. de		mm
22Kcor/1116/2896.11-	do		200
	d		30
ACQUISITION	Apr	2800.552	2800.55200
1a	HE	1b	1.00
1c	1.741	1d	1.00
1e	15036	1f	not used
1g	15036	1h	not used
1i	15036	1j	not used
1k	15036	1l	not used
1m	15036	1n	not used
1o	15036	1p	not used
1q	15036	1r	not used
1s	15036	1t	not used
1u	15036	1v	not used
1w	15036	1x	not used
1y	15036	1z	not used
1aa	15036	1ab	not used
1ac	15036	1ad	not used
1ae	15036	1af	not used
1ag	15036	1ah	not used
1ai	15036	1aj	not used
1ak	15036	1al	not used
1am	15036	1an	not used
1ao	15036	1ap	not used
1aq	15036	1ar	not used
1as	15036	1at	not used
1au	15036	1av	not used
1aw	15036	1ax	not used
1ay	15036	1az	not used
1ba	15036	1bb	not used
1bc	15036	1bd	not used
1be	15036	1bf	not used
1bg	15036	1bh	not used
1bi	15036	1bj	not used
1bk	15036	1bl	not used
1bm	15036	1bn	not used
1bo	15036	1bp	not used
1bq	15036	1br	not used
1bs	15036	1bt	not used
1bu	15036	1bv	not used
1bw	15036	1bx	not used
1by	15036	1bz	not used
1ca	15036	1cb	not used
1cc	15036	1cd	not used
1ce	15036	1cf	not used
1cg	15036	1ch	not used
1ci	15036	1cj	not used
1ck	15036	1cl	not used
1cm	15036	1cn	not used
1co	15036	1cp	not used
1cq	15036	1cr	not used
1cs	15036	1ct	not used
1cu	15036	1cv	not used
1cw	15036	1cx	not used
1cy	15036	1cz	not used
1da	15036	1db	not used
1dc	15036	1dd	not used
1de	15036	1df	not used
1dg	15036	1dh	not used
1di	15036	1dj	not used
1dk	15036	1dl	not used
1dm	15036	1dn	not used
1do	15036	1dp	not used
1dq	15036	1dr	not used
1ds	15036	1dt	not used
1du	15036	1dv	not used
1dw	15036	1dx	not used
1dy	15036	1dz	not used
1ea	15036	1eb	not used
1ec	15036	1ed	not used
1ee	15036	1ef	not used
1eg	15036	1eh	not used
1ei	15036	1ej	not used
1ek	15036	1el	not used
1em	15036	1en	not used
1eo	15036	1ep	not used
1eq	15036	1er	not used
1es	15036	1et	not used
1eu	15036	1ev	not used
1ew	15036	1ex	not used
1ey	15036	1ez	not used
1fa	15036	1fb	not used
1fc	15036	1fd	not used
1fe	15036	1ff	not used
1fg	15036	1fh	not used
1fi	15036	1fi	not used
1fk	15036	1fl	not used
1fl	15036	1fm	not used
1fo	15036	1fn	not used
1fp	15036	1fr	not used
1fq	15036	1ft	not used
1fr	15036	1fu	not used
1fs	15036	1fv	not used
1ft			

JK1457 Scan 2 (Av 2-6 Acq) 100%=8807 mv 8 Sep 92 17:20
LRP +EI SDJ 59/15F1 SP-19500 B.ro -FAB/MB, 381/1457, ZJY




Fig. 3

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/09623

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/195.1; 514/25, 75, 78; 536/4.1, 17.2, 17.9; 546/1, 292; 548/400; 549/5, 13, 29; 558/70, 88, 156

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS ONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X — A	Tetrahedron Letters, Volume 37, issued 1979, Van Boeckel et al, "Synthesis of glucosyl phosphatidylglycerol via a phosphotriester intermediate", pages 3561-3564, especially page 3563.	5-7, 12-14 and 17-18 — 1-4, 8-10 and 15-16
X — A	US, A, 4,684,625 (Eppstein et al) 04 August 1987, col. 8, lines 26-33.	5-7 — 1-4, 8-10 and 12-18



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be part of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

14 December 1993

Date of mailing of the international search report

28 DEC 1993

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. NOT APPLICABLE

Authorized officer

ELLI PESELEV

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US93/09623

A. CLASSIFICATION OF SUBJECT MATTER:
IPC (5):

A61K 9/00, 31/66, 31/70, 35/78; C07D 207/00, 327/00, 333/02, 335/00; C07F 9/02; C07H 15/00, 17/00

A. CLASSIFICATION OF SUBJECT MATTER:
US CL :

424/195.1; 514/25, 75, 78; 536/4.1, 17.2, 17.9; 546/1, 292; 548/400; 549/5, 13, 29; 558/70, 88, 156

THIS PAGE BLANK (USPTO)